

Research Article

The Influence of Patient Characteristics on Healthcare-Seeking Behavior: A Multilevel Analysis of 70 Primary Care Practices in Urban-Suburban Regions in Malta

Glorianne Pullicino

Visiting Assistant Lecturer, Department of Family Medicine, University of Malta, Specialist in Family Medicine, Primary Health Care Department, Malta

Philip Sciortino

Specialist in Family Medicine, Head of Department and Visiting Senior Lecturer, Department of Family Medicine, University of Malta, Malta

Liberato Camilleri

Head of Department and Associate Professor, Department of Statistics and Operations Research, University of Malta, Malta

Willemijn Schäfer

Researcher, International Comparative Health Services Research Primary care, Netherlands Institute for Health Services Research, Malta

Wienke Boerma

Senior Researcher and Consultant, Netherlands Institute for Health Services Research, Malta

ABSTRACT

Background: Social homogeneity and an almost indiscernible rural-urban difference are generally assumed to be strong factors that reduce any tendency for health inequities in a small island community. A strong primary health care system is one of the components that protect populations against inequities.

Aim: The aim of this study was to examine healthcare-seeking behaviour in urban and suburban regions in Malta.

Methods: The dataset of the Maltese arm of the QUALICOPC Project was analysed. A descriptive, cross-sectional study was designed. Seventy practicing general practitioners were selected randomly from the Malta Medical Council Family Medicine register after systematically removing the inactive practices. Ten patients presenting quasi-randomly in each primary care clinic completed a self-administered questionnaire. The chi-square test was used to test for differences in demographic and health care characteristics between the urban and suburban primary health care service.

Generalized Linear and Latent Mixed Models (GLLAMM) were used to perform the multilevel analysis using Stata/SE version 12.

Results: None of the 4 predictors (patients' primary or secondary educational level, age and gender) emerged to be significant for coping better with illness after GP visit. 82% of the total variance in this response (Yes/No) was between patients, 13% was between clinics and 5% between regions. General practitioners offered more health promotion services and asked more about polypharmacy when their patients were older.

Conclusion: Such findings provide information for policy makers to improve equity and resource allocations within the setting of urban Malta to help improve patients' outcomes, particularly for the at-risk or vulnerable population.

Keywords: Primary health care; Public health; Quality of care; Health care organization and management

How this fits in with quality in primary care?

What do we know?

Primary care is context-specific and complex with several inputs, processes and outputs. Multilevel modelling offers opportunities to explore primary care contextual effects and differences across regions.

What does this paper add?

We analyze and discuss multilevel models to demonstrate how patients' outcomes differ from clinic to clinic as well as between urban and suburban settings. The emergent data would be useful to inform policy makers, clinicians and health service researchers who aim to strengthen the primary health care system.

included important aspects regarding the process and outcomes of care, was defined. For the patient questionnaire the framework was based on the Consumer Quality Index of GP care.²¹

A pilot study was held in three countries in Europe (Belgium, the Netherlands and Slovenia) among a small sample of GPs and patients. GPs and patients were surveyed in the GP practice setting.⁹ The English version of the QUALICOPC questionnaire was translated into Maltese. A first translation was organized by a small team of people who were familiar with the PC practice in Malta and had a good understanding of English. A professional translator made an independent back-translation. Peer review indicated accepted face validity. Moreover, instrument development was further informed by focus groups. The researcher and the professional translator agreed on the final translated version. A pilot study was performed amongst fifty PC patients who did not form part of the sample. Further analysis indicated good test-retest reliability in both versions of the questionnaire.

Statistical Analysis

The chi-square test was used to test for differences in demographic and health care characteristics between the urban and suburban primary health care service. The data was analysed using the Statistical Package for Social Sciences (SPSS) version 20. Generalized Linear and Latent Mixed Models (GLLAMM) were used to perform the multilevel analysis using Stata/SE version 12.

Results

Seventy-eight GPs were invited to participate in this study. Eight of the latter did not consent to participate yielding an overall response rate of 88.6%. The response rates amongst the public and the private GPs were 94.3% and 82.9% respectively. A small remuneration was offered to the GPs. Amongst the private GPs who refused, four claimed that they were too busy to participate; another said that his patients would be very busy and another doctor claimed that his private clinic was too small for the fieldwork to occur. A public GP claimed that he was not interested in participating whilst another public GP was on long vacation leave during the time of the fieldwork. The patients' response rate in this study was 73%. Reasons for patients' non-participation included literacy issues, being too busy, disinterested or impatient.

The majority of the participants were females (61.9%, $n=386$). The sample population had an age distribution of 18-88 years with a mean of 48.2 years \pm 17.6 years. 53% of patients ($n=332$) visited GP clinics in urban areas whereas 47% ($n=294$) attended PC clinics in suburban villages. The highest level of education completed by the majority of participants was secondary education (45.1%, $n=282$) while tertiary education was completed by 36% ($n=225$) of the participants. Table 1 shows the socio-demographic and healthcare service characteristics among primary care patients in urban and suburban clinics.

Multilevel models showed that the effects of physician-level activities differed amongst clinics as well as between urban and suburban settings. Patients were asked whether in past 2 years, a GP from the practice had asked them about all the medications. 82% of the total variance in the responses (Yes/No) is between

Table 1: Socio-demographic and healthcare service patients' characteristics in urban and suburban primary care clinics.

Patients' Characteristics	Urban (n=332)		Suburban (n=294)	
	No.	%	No.	%
Gender				
Female	191	57.5	195	66.3
Male	140	42.2	98	33.3
Level of Education				
Never went to school/ Primary Education	65	55.1	53	44.9
Secondary Education	143	50.7	139	49.3
Postsecondary/Tertiary (E.g.: University)	123	54.7	102	45.3
Age				
15-29	63	19	47	16
30-44	80	24.1	86	29.3
45-59	88	26.5	81	27.6
60-74	75	22.6	57	19.4
75-89	26	7.8	23	7.8
Patient's Experiences and Views				
In past 2 years, a GP from the practice had asked about all the medications	105	50.5	103	49.5
In the past 12 months, a GP had talked about how to stay healthy	95	52.5	86	47.5
Patients were able to cope better with their health problem or illness after the GP visit	266	52.1	245	47.9

patients and 18% is between clinics. There was no variation between urban and suburban regions. Since the parameter estimate for each was positive, this indicated that the probability that the patient replied yes increased with an increase in age.

$$\text{Log}(p/1-p) = -1.4995 + 0.0599 \text{ primeduc} - 0.3362 \text{ seceduc} - 0.1147 \text{ sexpat} + 0.0192 \text{ agepat}$$

Health promotion was assessed by asking patients whether in the past 12 months, a GP had talked to them about how to stay healthy. 88% of the total variance in the responses (Yes/No) was between patients, 12% was between clinics and 0% between regions. Since the regression coefficient for each was positive, it indicated that the probability that the patient answered yes increased with an increase in age.

$$\text{Log}(p/1-p) = -2.5912 + 0.0041 \text{ primeduc} - 0.0009 \text{ seceduc} - 0.0420 \text{ sexpat} + 0.0332 \text{ agepat}$$

Patients' outcome was assessed by asking them whether

they were able to cope better with their health problem or illness after the GP visit. None of the 4 predictors (patient's educational status, sex and age) emerged to be significant since their p-value exceeded the 0.05 level of significance. 82% of the total variance in the responses (Yes/No) was between patients, 13% was between clinics and 5% between regions.

$\text{Log}(p/1-p) = 2.7570 - 0.0702 \text{ primeduc} - 0.5101 \text{ seceduc} - 0.5628 \text{ sexpat} + 0.0204 \text{ agepat}$

Patients in suburban regions tended to be able to cope better with their illness after they visited their GP.

Discussion

International research showed a lack of health promotion by GPs in the elderly.²² Conversely, in the present study, older patients were more likely to receive health promotion advice. One may question whether there is the unmet need for such services in the younger population. It could also be that the younger patients regard themselves as being healthy and do not value or seek health promotion services. Nevertheless, there may still be a lack of these services in the elderly, as the need may be a lot higher in this population. This means that there may also be unmet needs amongst the elderly. PC services could be responding to patients with different age groups in sync with their demands.

Similarly, elderly patients were more likely to be asked about polypharmacy. Younger PC patients might not have chronic diseases since they might be managed in secondary care specialized clinics. Moreover, the GP could associate polypharmacy with old age. Challenges of polypharmacy in the PC patient population can be addressed by collaborating with a pharmacist who can review the patient's medical records, make recommendations about low priority drugs or simplify regimes, for example, using long-acting medications, so that all tablets can be taken in the morning.²³

Rurality can contribute to the vulnerability of patients with chronic diseases through geographical barriers, financial constraints and limited availability of healthcare professionals.^{24,25} Moreover, patients may feel culturally marginalized in the urban health care context particularly if health literacy is low.²⁴ However, in the current study, patients in suburban regions tended to be able to cope better with their illness after they visited their GP. Qualitative systematic review and meta-synthesis postulated that rural communities supported long-term mutual doctor-patient relationships, feelings of a sense of belonging, self-reliant behavior and adaptation.^{24,25} In suburban Malta, it is unlikely for weather conditions or geographic distance to pose major barriers to access. These factors might mitigate feelings of vulnerability in suburban regions.²⁴

Potential limitations were identified in this study. Due to time and resource constraints, PC activities occurring in private hospitals and during home visits were excluded from this study. This study did not capture general practice activities carried out by other healthcare professionals, hospital-based specialists, private independent community-based specialists and unregistered medical practitioners. Participation and recall biases could have occurred. Participants could have tried to

answer in the way they believed the researcher or their GP wanted them to answer, rather than according to their own beliefs (the 'halo effect'). Respondents could have altered their responses in reaction to the researcher's presence or to the realization that they were being studied (the 'Hawthorne effect'). A training session was delivered to all fieldworkers to try to limit the inter-observer bias.

In view of the fact that this was a cross-sectional study, it was not able to demonstrate causality. There might have been an overrepresentation of patients who are frequent users of PC services. The reported improvement in the suburban areas might be the result of the service responding to the GPs' or the patients' demands rather than needs. Moreover, need might have been unperceived by patients, for example, in mental illnesses, and therefore, might not have led to demand. This study did not assess whether the services in urban and suburban areas were cost-effective and grounded in evidence-based medicine. Future quantitative research can address these limitations to ameliorate resource allocation and use, achieve value for money and improve patient outcomes.

Conclusion

In our study, we have analyzed a data set related to PC patients' experiences using multilevel modelling, which overcomes the limitations of normal regression models. Its findings would be useful to inform PC physicians, policy makers and health service researchers who aim to strengthen the primary health care system.

ETHICAL APPROVALS

The study was approved by the University Research Ethics Committee and the Data Protection Officer of the Primary Health Care Department.

ACKNOWLEDGEMENT

The Malta College of Family Doctors and the Malta Foundation Programme supported this project.

SOURCE OF FUNDING

This work was co-financed by the European Commission in the 7th Framework Programme under the Grant Agreement Number 242141.

REFERENCES

1. Brook RH, McGlynn EA, Cleary PD. Quality of health care Part 2: measuring quality of care. *N Engl J Med* 1996; 335: 966–970.
2. Shaw CD, Kalo I. A background for national quality policies in health systems 2002.
3. Legido-Quigley H, Dee MM, Nolte E, Glinos IA. Assuring the Quality of Health Care in the European Union. A case for action 2008.
4. Delnoij DMJ, Asbroek ten G, Arah OA, de Koning JS, Stam P, et al. Made in the USA: The import of American Consumer Assessment of Health Plan Surveys (CAHPS) into the Dutch social insurance system. *Eur J Public Health*. The Oxford University Press 2006; 16: 652–659.

5. Østerbye T, Sevaldsen J, Hansen KS, Freil M. Patients' experiences in Danish hospitals 2006.
6. Stubbe JH, Gelsema T, Delnoij DMJ. The Consumer Quality Index Hip Knee Questionnaire measuring patients' experiences with quality of care after a total hip or knee arthroplasty. *BMC Health Serv Res. BioMed Central Ltd* 2007; 7: 60.
7. Redding D, Boyd J. Data briefing. Involvement leads to satisfaction. *Health Serv J* 2008; 23: 17.
8. de Boer D, Delnoij D, Rademakers J. The discriminative power of patient experience surveys. *BMC Health Serv Res. BioMed Central Ltd* 2011; 11: 332.
9. Schäfer WLA, Boerma WG, Kringos DS, De Maeseneer J, Gress S, et al. QUALICOPC, a multi-country study evaluating quality, costs and equity in primary care. *BMC Fam Pract* 2011; 12: 115.
10. Agius F. Health and social inequities in Malta. *Soc Sci Med* 1990; 31: 313-318.
11. Statistics COO. Demographic review of the Maltese Islands 1985.
12. Turrell G, Kavanagh A, Subramanian SV. Area variation in mortality in Tasmania (Australia): the contributions of socioeconomic disadvantage, social capital and geographic remoteness. *Health Place*. 2006; 12: 291-305.
13. Hall SA, Kaufman JS, Ricketts TC. Defining urban and rural areas in U.S. epidemiologic studies. *J Urban Health. Springer US* 2006; 83: 162-175.
14. Prasad AM, Chakraborty G, Yadav SS, Bhatia S. Addressing the social determinants of health through health system strengthening and inter-sectoral convergence: the case of the Indian National Rural Health Mission. *Glob Health Action. Co-Action Publishing* 2013; 6: 1-11.
15. Gavos T. Health inequalities due to social determinants highlighted in international report 2013.
16. Zannoupas G, Lamnisos D, Kolokotroni O, Yiallourous P, Middleton N. Small-area mapping of premature mortality, rurality and deprivation indices on the small island of Cyprus. *The Oxford University Press* 2013; 23: 123-149.
17. EUROSTAT. Regions in the European Union. 2013; 1-144.
18. Pullicino G, Sciortino P, Calleja N, Schäfer W, Boerma W, Groenewegen P. Comparison of patients' experiences in public and private primary care clinics in Malta. *Eur J Public Health. The Oxford University Press* 2015; 25: 399-401.
19. Dickinson LM, Basu A. Multilevel modeling and practice-based research. *Ann Fam Med. American Academy of Family Physicians* 2005; 1: S52-S60.
20. Patterson L, Heller R, Robinson J, Birt C, van Ameijden E, et al. Developing a European urban health indicator system: results of EURO-URHIS 1. *Eur J Public Health* 2015 .
21. Schäfer WLA, Boerma WG, Kringos DS, De Ryck E, Gress S, et al. Measures of quality, costs and equity in primary health care instruments developed to analyse and compare primary care in 35 countries. *Qual Prim Care* 2013; 21: 67-79.
22. Badertscher N, Rossi PO, Rieder A, Herter-Clavel C, Rosemann T, et al. Attitudes, barriers and facilitators for health promotion in the elderly in primary care. A qualitative focus group study. *Swiss Med Wkly* 2012; 142: w13606.
23. Man M-S, Chaplin K, Mann C, Bower P, Brookes S, et al. Improving the management of multimorbidity in general practice: protocol of a cluster randomised controlled trial (The 3D Study). *British Medical Journal Publishing Group* 2016; 6: e011261.
24. Brundisini F, Giacomini M, DeJean D, Vanstone M, Winsor S, et al. Chronic disease patients' experiences with accessing health care in rural and remote areas: a systematic review and qualitative meta-synthesis. *Ont Health Technol Assess Ser. Health Quality Ontario* 2013; 13: 1-33.
25. Page-Carruth A, Windsor C, Clark M. Rural self-reliance: the impact on health experiences of people living with type II diabetes in rural Queensland, Australia. *Int J Qual Stud Health Well-being. Co-Action Publishing* 2014; 9: 24182.

ADDRESS FOR CORRESPONDENCE:

Dr. Glorianne Pullicino, MD MSc MRCP (Int) MMCDF, Visiting Assistant Lecturer, Department of Family Medicine, University of Malta, Specialist in Family Medicine, Primary Health Care Department, 36, 'Roma', Triq l-Alwetta. Mosta. MST4502, Malta, Tel: 79252218, E-mail: glorianneb@gmail.com